



Functional, Developmental and Morphological Integration: The Case of the Head and Forelimb in Bipedal Hominins

Citation

Lieberman, Daniel E., Dennis M. Bramble, David A. Raichlen, and Katherine W. Whitcome. 2008. Functional, developmental and morphological integration: The case of the head and forelimb in bipedal hominins. *American Journal of Physical Anthropology* 135(S46): 140-141.

Published Version

<http://dx.doi.org/10.1002/ajpa.20806>

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:2894773>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Functional, developmental and morphological integration: The case of the head and forelimb in bipedal hominins

Daniel E. Lieberman, Dennis M. Bramble, David A. Raichlen, Katherine W. Whitcome

Abstract:

Integration, a fundamental property of organisms, occurs via multiple mechanisms and for diverse reasons. Although there has been substantial work on the genetic and epigenetic mechanisms by which developmental integration occurs, we have less of an understanding of the evolutionary relationships between functional and developmental integration. In this respect, human evolution provides an interesting test case. In quadrupedal mammals, there is considerable functional integration among and between the limbs, but less integration between the limbs and the head. The evolution of bipedalism in hominins, however, provided a new opportunities for novel forms of integration by emancipating the forelimbs from any major role in locomotion. Here we consider how the forelimb and head become increasingly integrated in the genus *Homo* because of the biomechanical challenges of running. While the arm and the head interact little during walking, we have found that, during running, the stance side arm acts as a counterbalance to the head, stabilizing it against impulsive pitching forces generated by the heel strike transient. Moreover, the functional properties of this linkage may have driven several developmental changes in the proportions of the arm and the anatomy of the shoulder girdle during human evolution. Thus, evolutionary changes in arm and head morphology during human evolution may be more integrated than previously considered. This work was supported by funding from NFS.